

# Scritto d'esame di Elementi di Calcolo delle Variazioni

Pisa, 10 January 2017

1. Let us consider the functional

$$F(u) = \int_0^\pi (\dot{u}^2 - u \cos x) \, dx.$$

- (a) Discuss the minimum problem for  $F(u)$  with boundary condition  $u(\pi) = u(0)$ .
- (b) Discuss the minimum problem for  $F(u)$  with boundary condition  $u(\pi) = 2u(0)$ .

2. Let us consider the boundary value problem

$$\ddot{u} = \frac{\sinh u - 3}{3 + \cosh u}, \quad u(0) = u(2017) = 3.$$

- (a) Discuss existence, uniqueness, regularity of the solution.
- (b) Prove that the solution satisfies  $1 < u(x) \leq 3$  for every  $x \in [0, 2017]$ .

3. Let us consider, for every  $\ell > 0$ , the problem

$$\inf \left\{ \int_0^\ell (\dot{u}^2 + \arctan u - u^2) \, dx : u \in C^1([0, \ell]), \, u(0) = u(\ell) = 0 \right\}.$$

- (a) Determine for which values of  $\ell$  the infimum is negative.
- (b) Determine for which values of  $\ell$  the infimum is actually a minimum.

4. Let us set

$$m_\varepsilon = \min \left\{ \int_0^1 (\varepsilon \ddot{u}^2 + \cos \dot{u} + \cos u) \, dx : u \in C^2([0, 1]), \, u(0) = u'(0) = 1 \right\}.$$

- (a) Prove that  $m_\varepsilon$  is well-defined (namely the minimum exists) for every positive  $\varepsilon$ .
- (b) Compute the limit of  $m_\varepsilon$  as  $\varepsilon \rightarrow 0^+$ .

Every step has to be *reasonably* motivated. Every exercise is marked considering the *correctedness* of the arguments provided and the *clarity* of the presentation. Just writing the answer without explanations deserves no marks.